

Anxiety and Depression during pregnancy and the postnatal period



What the fact sheet covers:

- Types of depression during pregnancy and the postnatal period
- Symptoms of postnatal depression
- Symptoms of perinatal anxiety
- Causes of pregnancy-related depression
- Risk factors and triggers for pregnancy-related depression
- Finding help.

What is perinatal depression?

Perinatal depression and anxiety refer to changes in mood that may occur during pregnancy or the postnatal period. Approximately 15–20 per cent of women in Australia are affected by perinatal depression or anxiety. The childbearing years, especially the first few weeks after childbirth, are the peak period for the onset of depression in women. Perinatal depression varies with respect to symptoms, timing of onset, causes, risk factors, severity and duration. Perinatal mental health issues vary with respect to symptoms, timing of onset, causes, risk factors, severity and duration. It can also vary in the need for professional assessment and the type of treatment.

Types of depression during pregnancy and the postnatal period

Short episodes of tiredness, nausea, aches and pains, irritability, sleep disturbance and loss of interest in sex are relatively common as part of the normal adjustment process in the perinatal period and will not require treatment.

It is important to distinguish between the 'baby blues', and other perinatal mental health issues that warrant professional treatment.

1. The 'baby blues'

The term 'baby blues' refers to a brief episode of mood swings, tearfulness, anxiety and difficulty with sleeping that is very common in the first



week after the birth of a baby. It requires no special treatment, unless the symptoms are severe.

2. Antenatal depression

Antenatal depression refers to depression that starts during pregnancy. Between 10–15 per cent of pregnant women experience mood swings during pregnancy that last more than two weeks at a time and interfere with normal day-to-day functioning. Medical assessment is necessary in such circumstances.

3. Postnatal depression (PND)

PND describes the more severe or prolonged symptoms of depression (clinical depression) that last more than two weeks and interfere with the ability to function with normal routines on a daily basis including caring for a baby. Around 14 per cent of women in Australia (one in seven) experience PND. For around 40 per cent of these women, the symptoms begin in pregnancy.

4. Postnatal (puerperal) psychosis

This is an uncommon disorder that occurs in 1–2 individuals per 1,000 women. It has a sudden onset with severe symptoms – usually within two to three weeks of childbirth. Symptoms can also begin during pregnancy, especially where there has been a prior episode of psychosis or bipolar disorder. This illness requires urgent medical assessment and treatment.

Symptoms of PND

There can be differences in the nature, severity and duration of the symptoms of depression seen in women who are pregnant or have recently given birth. Professional help is generally required to diagnose the type of depression and decide the best approach to treatment.

Common symptoms of PND include:

- Loss of enjoyment in usual activities
- Loss of self-esteem and confidence
- Loss of appetite and weight, or weight gain
- Difficulty with sleep (irrespective of the baby's routine)

- A sense of hopelessness and of being a failure
- A wish not to be alive
- Suicidal thoughts or ideas
- Panic attacks
- Loss of libido
- Fears for the baby's or partner's safety or wellbeing.

It is also very important that any talk of suicide be taken seriously and treatment from a mental health professional or other appropriate person be immediately sought.

The symptoms of postnatal psychosis can be severe and include the following features:

- Confused thinking
- Hallucinations
- Restlessness, agitated behaviours, or strange movements
- Fearfulness and worrying (often about the baby)
- Mood swings, sometimes with inappropriate emotions
- Elevated mood, extremely heightened energy levels and manic behaviours
- Inability to sleep
- Appearing out of touch with reality (psychotic), being suspicious, or demonstrating inappropriate behaviours.
- Medical assessment is necessary if any of these symptoms are present.

Perinatal Anxiety

While pregnancy and the arrival of a new baby can be very exciting, most women experience some worries about things like having a healthy pregnancy, delivering the baby, keeping their baby safe, and potential impacts on their relationship, career, or finances. For some people, those worries can become overwhelming and unmanageable. Anxiety during pregnancy is often under-diagnosed and 10–20% of women will experience clinically significant anxiety during pregnancy or the postnatal period.





Symptoms of perinatal anxiety

Symptoms may include excessive worry, a sense of dread, physical symptoms of anxiety such as elevated breathing and heart rate, stomach distress, and difficulty sleeping.

Causes of pregnancy-related depression and anxiety

There are a variety of causes or triggers that can lead to the onset of clinical depression and anxiety during pregnancy and the postnatal period.

Some mood disorders such as puerperal psychosis and bipolar disorder are known to be linked to biological (genetic or biochemical) causes.

Perinatal depression and anxiety can also be associated with psychosocial stressors such as:

- Maladaptive behaviour patterns and dysfunctional thought processes and coping styles
- Problems with key relationships
- Difficult living conditions and negative life events.

Risk factors and triggers for pregnancy related depression and anxiety include:

- A previous history of depression, bipolar disorder or psychosis
- Stressful life events
- Lack of social supports

- A history of physical, sexual or emotional abuse
- Pregnancy loss
- Childbirth-related distress
- A baby that is difficult to settle, restless or unwell
- Personality types that increase vulnerability to depression such as the anxious worrier or socially avoidant personality styles.

Finding help

Various health professionals and allied health professionals are qualified to help people experiencing depression during pregnancy and the postnatal period including:

- Doctors – general practitioner (GP)
- Obstetrician
- Psychiatrist
- Midwives
- Child and family health nurses
- Social workers
- Counsellors
- Psychologists.

The [Gidget Foundation](#) offers a free video counselling service for some new and expecting parents who are experiencing perinatal depression and/or anxiety and are unable to have a face-to-face consultation.

Check your eligibility by contacting contact@gidgetfoundation.org.au or 1300 851 758.



**Black Dog
Institute**

Key points to remember

- The childbearing years, particularly the first few weeks after childbirth, are the peak period for onset of depression in women.
- Depression can begin during pregnancy.
- Excessive fatigue can contribute to low mood so adequate rest can help to prevent exhaustion.
- Around 14 percent of women will suffer postnatal depression.
- Anxiety and depression often go hand-in-hand.

For advice about medications during pregnancy and breastfeeding

- Speak to your specialist, GP or pharmacist for safety advice about using medications while pregnant or breastfeeding.

Crisis support

If you need support, call one of the following numbers:

- [Lifeline Australia](https://www.lifeline.org.au) – 13 11 14 (24/7)
- [PANDA](https://www.panda.org.au) [Perinatal Anxiety and Depression Australia] National Helpline (Mon to Fri, 9am– 7.30pm AEST) 1300 726 306



For more information

Visit our website blackdoginstitute.org.au
Find us on social media @blackdoginst



This document may be freely downloaded and distributed on condition no change is made to the content. The information in this document is not intended as a substitute for professional medical advice, diagnosis or treatment. Not to be used for commercial purposes and not to be hosted electronically outside of the Black Dog Institute website.



**Black Dog
Institute**